

Leukaemia Section

Short Communication

t(4;12)(q21;p13)

Tatiana Gindina

R.M.Gorbacheva Memorial Institute of Children Oncology, Hematology and Transplantation at First Saint-Petersburg State Medical University named I.P.Pavlov, Saint-Petersburg, Russia /
tatgindina@gmail.com

Published in Atlas Database: September 2018

Online updated version : <http://AtlasGeneticsOncology.org/Anomalies/t0412q21p13ID1504.html>

Printable original version : <http://documents.irevues.inist.fr/bitstream/handle/2042/70533/07-2018-t0412q21p13ID1504.pdf>

DOI: 10.4267/2042/70533

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2019 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

Review on t(4;12)(q21;p13) in lymphoid malignancies

Keywords

Chromosome 4; Chromosome 12; Acute lymphoblastic leukemia; Primary mediastinal large B-cell lymphoma; Adult T-cell leukemia/lymphoma; Acute megakaryoblastic leukemia.




Figure 1. Partial karyotype with t(4;12)(q21;p13) in a female patient with B-ALL (table 1, #8).

Clinics and pathology

Disease

Translocation t(4;12)(q21;p13) occurs predominantly in B-cell lymphoid disorders, including ALL, NHL, rarely with T-ALL and

AML.

Acute lymphoblastic leukemia (ALL): was diagnosed in 5 patients (Groupe Français de Cytogénétique Hématologique, 1993; Behm et al., 1996; Elghezal et al., 2001; Gindina T., own case, table 1, #8).

Primary mediastinal large-B-cell lymphoma was found in 1 patient (Palanisamy et al, 2002)

Adult T-cell Leukemia: 1 patient (Sadamori et al., 1991).

Non-Hodgkins lymphoma: 1 patient (Schouten et al., 1990).

Acute myeloid leukemia: 1 patient with AML-M7 (Ohyashiki et al., 1984).

Phenotype/cell stem origin

Two patients had B-cell Early B ALL (CD10+) and Pre-B ALL (CIG +) (Groupe Français de Cytogénétique Hématologique, 1993).

Epidemiology

The translocation t(4;12)(q21;p13) was found in adults and children as well as equally among male and female patients.

Genetics

Genes implicated in this translocation remain unknown.

Pts	Disease	Gender/Age	Karyotype
1	NHL	F/?	49,XX, +3,+18,t(3;?)(q29;?), t(4;12)(q21;p13) , del(6)(q23),t(7;?)(q21;?),t(9;?)(p23;?),t(14;18)(q32;q21),t(17;?)(p13;?),+mar
2	ATL	M/?	48,XY,+del(1)(p32),+3, t(4;12)(q21;p13)
3	PMBL	F/?	50,XX, dup(2)(q21q31),+3, t(4;12)(q21;p13) , del(6)(q23),del(7)(q21),add(9)(p23),add(13)(p13),t(14;18)(q32;q21),+18, +add(20)(p13),+21
4	B-ALL	F/?	47,XX, t(4;12)(q13;p12) ,del(15)(q14q25),+21,der(21) t(12;21)[3]/ 48,idem,+13[1]/49,idem,+X[1]
5	B-ALL Remission duration 30+	F/6	47,XX,t(1;12)(p22;p13), t(4;12)(q21;p13) ,+10, del(11)(q23)/45,X,-X,t(4;12)
6	B-ALL	M/3	47,XY,del(1q),del(2q),der(3q), t(4;12)(q13;p12) , del(6)(q21q25), + mar
7	B-ALL	M/5	46, XY, der(1)t(1;?)(p36;?), t(4;12)(q13;p12)
8	B-ALL	F/3	48,XX, t(4;12)(q21;p13) ,+10,+21
9	AML-M7	M/59	46,XY, t(4;12)(q21;p13)

Table 1. Reported cases with t(4;12)(q21;p13).

NHL: non-Hodgkin's lymphoma; ATL: Adult T-cell leukemia; PMBL: Primary mediastinal large-B-cell lymphoma; B-ALL: B-Acute lymphoblastic leukemia; AML-M7: Acute megakaryoblastic leukemia

1. Schouten et al., 1990; 2. Sadamori et al., 1991; 3. Palanisamy et al., 2002; 4. Elghezal et al., 2001; 5. Behm et al., 1996; 6,7. Groupe Français de Cytogénétique Hématologique, 1993; 8. Gindina, personal observation; 9. Ohyashiki et al., 1984.

Cytogenetics

Additional anomalies

Additional chromosome abnormalities were observed in all these patients with lymphoid malignancies. Translocation t(4;12) was associated with such anomaly as translocation t(14;18)(q32;q21) in 2 cases (Schouten et al., 1990; Palanisamy et al., 2002) and translocation t(12;21)(p13;q22) ETV6/ RUNX1 in 1 patient (Elghezal et al., 2001).

Trisomies of chromosomes 3 and 21 were present in 3 patients each (Schouten et al., 1990; Sadamori et al., 1991; Palanisamy et al., 2002; and Palanisamy et al., 2002; Elghezal et al., 2001; Gindina T. own case #8, respectively), trisomies of chromosomes 10 and 18 were in 2 cases each (Behm et al., 1996; Gindina T. own case #8, respectively and Schouten et al., 1990; Palanisamy et al., 2002, respectively). Deletion of 6q was demonstrated in 3 patients (Schouten et al., 1990; Palanisamy et al., 2002; Groupe Français de Cytogénétique Hématologique, 1993). In most cases, t(4;12)(q21;p13) was as a part of a complex karyotype. As a single anomaly, t(4;12)(q21;p13) was described only in one case of AML (Ohyashiki et al., 1984).

Result of the chromosomal anomaly

Fusion protein

Oncogenesis

Most likely, the translocation t(4;12)(q21;p13) is a secondary genetic event in oncogenesis.

References

Behm FG, Raimondi SC, Frestedt JL, Liu Q, Crist WM, Downing JR, Rivera GK, Kersey JH, Pui CH. Rearrangement of the MLL gene confers a poor prognosis in childhood acute lymphoblastic leukemia, regardless of presenting age. *Blood*. 1996 Apr 1;87(7):2870-7

Collaborative study of karyotypes in childhood acute lymphoblastic leukemias. Groupe Français de Cytogénétique Hématologique. *Leukemia*. 1993 Jan;7(1):10-9

Elghezal H, Le Guyader G, Radford-Weiss I, Perot C, Van Den Akker J, Eydoux P, Vekemans M, Romana SP. Reassessment of childhood B-lineage lymphoblastic leukemia karyotypes using spectral analysis. *Genes Chromosomes Cancer*. 2001 Apr;30(4):383-92

Ohyashiki K. Nonrandom cytogenetic changes in human acute leukemia and their clinical implications *Cancer Genet Cytogenet* 1984 Apr;11(4):453-71

Palanisamy N, Abou-Elella AA, Chaganti SR, Houldsworth J, Offit K, Louie DC, Terayu-Feldstein J, Cigudosa JC, Rao PH, Sanger WG, Weisenburger DD, Chaganti RS. Similar patterns of genomic alterations characterize primary mediastinal large-B-cell lymphoma and diffuse large-B-cell lymphoma *Genes Chromosomes Cancer* 2002 Feb;33(2):114-22

Sadamori N, Isobe M, Shimizu S, Yamamori T, Itoyama T, Ikeda S, Yamada Y, Ichimaru M. Relationship between chromosomal breakpoint and molecular rearrangement of T-cell antigen receptors in adult T-cell leukaemia *Acta Haematol* 1991;86(1):14-9

Schouten HC, Sanger WG, Weisenburger DD, Armitage JO. Abnormalities involving chromosome 6 in newly diagnosed patients with non-Hodgkin's lymphoma Nebraska Lymphoma Study Group *Cancer Genet Cytogenet*

This article should be referenced as such:

Gindina T. t(4;12)(q21;p13). *Atlas Genet Cytogenet Oncol Haematol*. 2019; 23(9):293-294.